

SUPERCURRENT STATES IN 1D FINITE-SIZE RINGS

Vladimir A. Kashurnikov and Alexei I. Podlivaev

Moscow State Engineering Physics Institute, 115409 Moscow, Russia

Nikolai V. Prokof'ev and Boris V. Svistunov

Russian Research Center "Kurchatov Institute", 123182 Moscow, Russia

Abstract

We consider topological supercurrent excitations (SC) in 1D mesoscopic rings. Under certain conditions such excitations are well-defined except for (i) a tunneling between resonating states with clockwise and anti-clockwise currents, which may be characterized by the amplitude Δ , and (ii) a decay of SC assisted by phonons of the substrate, both effects being macroscopically small. Our approach being based on the hydrodynamical action for the phase field and its generalization to the effective Hamiltonian explicitly takes into account transitions between the states with different topological numbers and turns out to be very effective for the calculation of Δ and estimation of the decay width of SC, as well as for the unified description of all known 1D superfluid-insulator transitions.

Most attention is paid to the calculation of the macroscopic scaling of Δ (the main superfluid characteristic of a mesoscopic system) under different conditions: a commensurate system, a system with single impurity, and a disordered system. The results are in a very good agreement with the exact-diagonalization spectra of the boson Hubbard models.

Apart from really 1D electron wires we discuss two other important experimental systems: the 2D electron gas in the FQHE state and quasi-1D

superconducting rings. We suggest some experimental setups for studying SC, e.g., via persistent current measurements, resonant electro-magnetic absorption or echo signals, and relaxation of the metastable current states.

I. INTRODUCTION

The problem we address here is a general one for the condensed matter physics and arises each time when we deal with the sample of nontrivial topology. Namely, if we write down the field operator in terms of density and phase $\Psi(x) = |\rho(x)|^{1/2} e^{i\Phi(x)}$, then in a sample with the topology of a ring we may define the topological excitations characterized by the winding number

$$I = \frac{1}{2\pi} \oint_{\Gamma} \nabla \Phi d\vec{l}, \quad (1)$$

where the integration contour Γ is around the ring hole. Clearly, these excitations exist in any space dimension and for $I \neq 0$ carry current due to nonzero gradient of the phase field. The stability of topologically excited states does depend on the dimensionality and properties of the bulk system. In the normal state of the 3D and 2D liquid the relaxation time of the SC due to disorder or crystal potential is extremely short (of order of the transport scattering rate). However below the superconducting or superfluid phase transition the relaxation mechanism of I is only through the vortices, which are macroscopic objects in a sense that their energy is system size dependent (the pairs of vortices have finite energy, but their only effect on I is to provide short-time fluctuations in I leaving the long-time average of the topological quantum number unmodified). In the ground state or at a very low temperature the probability of vortex creation by quantum or thermal fluctuations can be ignored, and the integral (1) is a true invariant of the system. Note also, that at finite temperatures below the critical point there may be introduced a regularized operator of phase in terms of which I is also an exact (with macroscopic accuracy) constant of motion [1].

The above argument does not work in 1D, because one can not define vortex in one spatial dimension. It means that the dynamics of I is defined by microscopic parameters

of the problem, and the bare transition amplitude between different quantum numbers is independent of the system size. Calculating the relaxation of the topological excitations using bare parameters of the Hamiltonian we conclude in agreement with the conventional wisdom that disorder or periodic potential does not allow conserving current states in 1D. The crucial point for the study presented here is that the SC in the interacting 1D system are always "dressed" by density fluctuations. Any attempt to change I would result in driving the wavefunction of the whole ring into orthogonal state (that is $\langle I | I' \neq I \rangle \rightarrow 0$ with macroscopic accuracy - see also below). Now, the renormalized amplitude connecting different numbers I is size-dependent for the lowest excited states. Depending on the interaction parameters between the particles in the original model, one can obtain arbitrarily long relaxation times for the topological excitations.

In this paper we discuss the stability and relaxation of SC in 1D interacting systems. To be definite, most of the time we consider Bose systems. As is well known [2], the statistics and interaction effects are indistinguishable in 1D, so that fermions are exactly described as hard-core bosons, and spin chains are converted into the interacting bosons by Holstein-Primakoff transformation. Our main interest is in the low-energy excitation spectrum of the ring in the superfluid phase, including the energy splittings between the resonating SC and their hybridization with the sound modes.

Experimentally SC may be observed in the response of the ring to the vector potential. Suppose that at the initial moment of time the system is in the ground state with some value of $I = I_G$. In the general case I_G is a function of the magnetic flux through the ring and the parity of the total number of particles in the system: for bosons and odd parity fermion systems in zero flux $I_G = 0$; fermion ring with the even number of particles is equivalent to a bosonic system with the half-quantum of the magnetic flux through the ring. In this case the ground state is degenerate between $I = 1$ and $I = 0$. By switching on instantaneously an integer number of flux quanta, $\phi = m\phi_0 = mhc/e$, we excite nonadiabatically the supercurrent state with $I = I_G + m$. One may then study the time evolution of I by measuring e.g. the oscillations and decay of the magnetic moment generated by the current. In a static

experimental setup the relevant quantity is the magnitude and sign of the persistent current near the points of the ground state degeneracy. In bosonic and odd parity fermionic systems these points are given by $\phi \approx (m + 1/2)\phi_0$. Near these points the persistent current has the maximum possible amplitude and abruptly changes its sign. The width of the crossover region is very small $|\phi - (m + 1/2)\phi_0| \ll \phi_0$ and directly related to the frequency of the current oscillations in the dynamic experiment. In the last section we will describe in some detail the most promising experimental 1D system - edge states of the 2D electron gas in the regime of fractional quantum Hall effect (FQHE). Here the SC are related to the electrical charge transfer between the edges in the Carbino disk geometry thus making it possible to study the phenomenon by monitoring the charging effects between the edges. Finally, we will demonstrate that the physics of SC relaxation in 1D systems can be directly applied to a 3-dimensional superconducting ring with the ring length being much longer than the wire cross-section diameter.

The other way to excite the SC is by pumping the system with the high-frequency electromagnetic radiation in resonance with the energy of, say, the first current state.

The paper is organized as follows. In section II we formulate the effective-action approach to the problem and show it to be very convenient for unified description of the critical points and renormalization group (RG) equations for the superfluid-insulator transitions in commensurate and disordered systems. Next we calculate the energy splitting, Δ_{AB} , between degenerate current groundstates (in the presence of the half-quantum of magnetic flux or its equivalent) using an instanton technique. This splitting and its dependence on the ring length L is the main mesoscopic characteristic of SC. It gives the frequency of the coherent current oscillations in the superfluid phase, while its divergence signals about the superfluid-insulator transition. All calculations in section II are done with the logarithmic accuracy.

In section III we formulate an alternative approach based on the effective Hamiltonian for the SC dynamics. This way we calculate precisely the hybridization coupling between the topological states and phonons, which allows to fix all the ratios between the lowest

energy levels. The other goal is the possibility to account for the resonances between the SC and sound waves and describe correctly level crossings. All the results obtained analytically in sections II and III are tested in section IV by intensive numeric calculations done on finite-size Hubbard-like ring models. We present the low-energy spectra for various systems and discuss the procedure of their extrapolation on larger system size by combining the numeric data with the analytic RG equations. In section V we discuss possible experimental setups for the study of SC. We also considered in some detail two important experimental systems, i.e., the 2D electrons in the FQHE state and a long, narrow 3D superconducting ring, as the most promising systems where our results may be tested.

II. EFFECTIVE-ACTION APPROACH

Since the stability of the SC states is associated with the conservation of the topological invariant I (1) the most natural language for the description of its violation or even complete disappearance under various conditions is that which can directly take into account processes changing the value of I . In this section we introduce such a language in terms of the effective action for the phase field in (1+1) dimensions.

Consider Popov's hydrodynamical action [3] with the density-field fluctuations integrated out:

$$S[\Phi] = \int dx d\tau \left\{ i n_0(x) \Phi'_\tau + \frac{1}{2} \Lambda_s (\Phi'_x)^2 + \frac{1}{2} \kappa (\Phi'_\tau)^2 \right\} . \quad (2)$$

Here $\Phi(x, \tau)$ is the phase field (x is the spatial coordinate and τ is the imaginary time), $n_0(x)$ is the local number density with the coordinate dependence coming from this or that short-range external potential, Λ_s is the superfluid stiffness, and $\kappa = (d\mu/dn)^{-1}$ is the compressibility (μ is the chemical potential and n is the macroscopic density). As we shall see below, the dependence of n_0 on x in (2) is of prime importance being responsible for a particular type of possible violation of superfluidity. So $n_0(x)$ can not be replaced by n .

Zero-point fluctuations with the change of I are represented by the topological defects (vortices) in the field $\Phi(x, \tau)$. Hence, the question of the existence in a ring system of the

metastable groundstate-like excitations with nonzero expectation values of I is the question of statistics of the vortices. If in the long-range limit the vortices are absent (bounded in short-range pairs) such states are possible. If separate vortices are present at the arbitrarily large scales one can not introduce the topological quantum number and the only possible equilibrium expectation value of I is zero. In this case the very description of the system in terms of the effective action (2) is, generally speaking, no longer valid because of divergent long-range renormalizations.

The self-consistent criteria for superfluidity under different conditions (commensurate external potential, disorder, or a special case with the single impurity) follow directly from the action (2). With rescaled variable τ this action reads

$$S[\Phi] = \int dx dy \left\{ i n_0(x) \Phi'_y + \frac{1}{2\pi K} (\nabla \Phi)^2 \right\} . \quad (3)$$

Here $y = c\tau$ ($c = \sqrt{\Lambda_s/\kappa}$ is the sound velocity),

$$K^{-1} = \pi \sqrt{\Lambda_s \kappa} . \quad (4)$$

The first term in (3) is of topological nature and "reacts" only on topological defects. Integrating over y and introducing

$$\gamma(x) = -2\pi \int_0^x n_0(x') dx' \quad (5)$$

one may rewrite $S[\Phi]$ as

$$S[\Phi] = \frac{1}{2\pi K} (\nabla \Phi)^2 + i \sum_j p_j \gamma(x_j) , \quad (6)$$

where j enumerates the vortices, $p_j = \pm 1, \pm 2, \dots$ and x_j are the charge and the x -coordinate of the vortex j , respectively. The effective action (6) has the form of XY -model (the first term) with the additional feature that each vortex brings in a spatially- and charge-dependent phase (the second term). This phase plays an extremely important role. To realize this, consider a homogeneous system. In the absence of the second term in the action (6) the system would be equivalent to the XY -model and thus would demonstrate

Kosterlitz-Thouless transition at $K = 1/2$ which means destruction of superfluidity. However, it is clear that the homogeneous system is always trivially "superfluid" because of the momentum conservation. Hence, no phase transition actually should occur, and it is the second term in Eq.(6) that corrects the result. Indeed, in the homogeneous case $\gamma(x_j)$ is a linear function of x_j and the integration over x_j makes the net contribution of the vortex j to the partition function to be zero. A similar picture takes place in a periodic incommensurate potential (and, in particular, in a lattice at an incommensurate filling). In this case $\gamma(x)$ may be represented as

$$\gamma(x) \equiv \gamma(s, \xi) = -2\pi\nu s - 2\pi \int_0^\xi n_0(x') dx' , \quad (7)$$

where $x = sl + \xi$, l is the period of the potential, s is the corresponding number of periods, $\xi \in [0, l]$ is the reduced coordinate, $\nu = \int_0^l n_0(x) dx \equiv nl$ is the filling factor. The second term in the r.h.s. of Eq.(7) leads only to a certain short-range renormalization of the statistical weight of the vortex and, thus, does not play an essential role, so we omit this term from now on. (Note also, that in a lattice this term is absent from the very beginning.) The first term in the r.h.s. of Eq.(7) is linear in s . This means that just as in the homogeneous case the net contribution of vortices to the partition function is zero, provided ν is not a rational number. Hence, no phase transition occurs.

In the case of commensurate filling the discreteness of s becomes important. Now there appears a class of vortices whose net contribution is nonzero. These are the vortices whose charges are the multiples of p , the denominator of the filling factor. All these vortices enter the partition function with one and the same spatially independent phase, so one deals with the XY-model with the constraint that the charge of a vortex be a multiple of p . Thus, the system should experience a Kosterlitz-Thouless-type phase transition at $K = p^2/2$, which can be identified with the known superfluid - Mott insulator transition in a commensurate system [4,2,5]. Note, that the critical value of K follows immediately from the effective action (6) by the free-energy-sign argument for a single vortex [6]. This argument is exact and need not a resorting to the RG, provided K is understood as the long-range asymptotic

quantity [1]. Nevertheless, RG equations are of interest for some applications (see below). They read (cf. [5])

$$\begin{cases} dK/d\lambda = w^2 \\ dw/d\lambda = (2 - p^2/K) w \end{cases} \quad (8)$$

Here $\lambda = \ln R$, $R = \sqrt{x^2 + y^2}$ being the characteristic scale of distance, $K(\lambda)$ is the mesoscopic value of K . Eqs.(8) are the standard Kosterlitz-Thouless RG equations [6,7] written for the vortices of the charge p . As usual, $(w(\lambda))^2$ is proportional to the number of vortex pairs of the size $\sim R$ in the area $\sim R^2$ and is much less than unity in the region of applicability of Eqs.(8). Mesoscopic values of Λ_s and κ are related to $K(\lambda)$ by

$$\Lambda_s(\lambda) = \frac{c}{\pi K(\lambda)}, \quad \kappa(\lambda) = \frac{1}{\pi c K(\lambda)}, \quad (9)$$

c being irrenormalizable quantity.

Considering disordered superfluid system it is convenient to rewrite the second term in (6) in terms of the pairs of vortices: $\sum_j p_j \gamma(x_j) \rightarrow \sum_\zeta q_\zeta \tilde{\gamma}(x_{1\zeta}, x_{2\zeta})$, where ζ indexes the vortex pairs, $q_\zeta = 1, 2, 3, \dots$ is the charge of the vortices in the pair ζ , $x_{1\zeta}$ and $x_{2\zeta}$ are the x -coordinates of the positive and the negative vortex in the pair ζ , respectively, and

$$\tilde{\gamma}(x_1, x_2) = -2\pi \int_{x_1}^{x_2} n_0(x) dx. \quad (10)$$

Being interested in the long-range contribution of a vortex pair we may average its phase factor over short-range translations along the x -axis:

$$e^{i2\pi q \tilde{\gamma}(x_1, x_2)} \rightarrow \langle e^{i2\pi q \tilde{\gamma}(x_1 + \xi, x_2 + \xi)} \rangle_\xi = f(x_1 - x_2). \quad (11)$$

Since $n_0(x)$ in a disordered system is a random function with some microscopic correlation length, the function $f(x_1 - x_2)$ decays at microscopically small $|x_1 - x_2|$ and may be replaced by $\propto \delta(x_1 - x_2)$. We see thus, that in a disordered system the second term in the effective action (6) is equivalent to the constraint that only "vertical" vortex pairs be present. This constraint changes the entropy per vortex as compared to the regular commensurate case. Now the entropy per vortex is $(3/2) \ln L + \text{finite term}$ since the entropy

of a pair is obviously $3 \ln L + \text{finite term}$. The energy of a vortex with the minimal charge is $K^{-1} \ln L + \text{finite term}$, so the free-energy-sign argument yields the critical macroscopic value of $K = 2/3$, in agreement with the result of Giamarchi and Schulz [5] obtained from other considerations. The renormalization group for the disordered case may be easily derived from the effective action in analogy with the Kosterlitz-Thouless treatment [6]. The new feature associated with the above-mentioned constraint is that renormalizable quantities now are Λ_s and c while κ can not be renormalized by the vertical vortex pairs. The RG equations read now

$$\begin{cases} dK/d\lambda = Kw^2 \\ dw/d\lambda = (3/2 - 1/K)w \end{cases} \quad (12)$$

Here λ and w have the same meaning as in Eqs.(8). Mesoscopic values of Λ_s and c are related to $K(\lambda)$ by

$$\Lambda_s(\lambda) = \frac{1}{\kappa(\pi K(\lambda))^2}, \quad c(\lambda) = \frac{1}{\pi \kappa K(\lambda)}. \quad (13)$$

Since in the disordered case there takes place a renormalization of c , a question may arise of what quantity c should be used for the introduction of the variable $y = c\tau$. For the derivation of Eqs.(12) it is natural to choose c to be some small-scale value of $c(\lambda)$. For the application of RG to a particular mesoscopic system the most appropriate choice is $c = c(\lambda_*)$, where λ_* is the large-scale cutoff for λ , related to the size of the system. We notice, however, that this difference leads only to small corrections in the parameter $w^2 \ll 1$ to the RG equations, and thus may be neglected.

Finally, consider a special case of a regular system with a single impurity (or any other defect). As it follows from the above analysis for regular and disordered systems, in this case the effective action corresponds to XY -model with the constraint that the vortex pairs be vertical and located only in the vicinity of the line $x = x_0$, x_0 being the coordinate of the defect. The free-energy-sign argument (entropy per vortex is $\ln L + \text{finite term}$) immediately yields the critical value of $K = 1$, in accordance with the result of Kane and Fisher [8]. In the case of one defect no renormalization of K occurs. Physically it is clear since the

transition is associated only with the renormalization of the strength of the defect: effective tunneling constant tends to zero at $K > 1$ [8].

Now we use the effective action to consider the splitting Δ_I of the SC level I in a finite-size system as a function of L and K . This splitting is due to the tunneling between the states $|I\rangle$ and $|-I\rangle$ which means that the true eigenstates of the finite-size system are superpositions of $|I\rangle$ and $|-I\rangle$. To calculate Δ_I *ab initio* one may take advantage of the conventional instanton approach starting from the formula

$$\langle -I | e^{iHt} | I \rangle = e^{-i(E_g + E_I)t} \sin(\Delta_I t), \quad (14)$$

where E_g is the groundstate energy, $E_I = 2\pi^2 I^2 \Lambda_s / L$ is the supercurrent energy, H is the system Hamiltonian. The next step is to use the imaginary time $t = -i\beta$ in order to represent the l.h.s. of (14) in terms of the effective action (2). Since the functional integral is naturally introduced in terms of the coherent states, we represent $|I\rangle$ as the expansion over the set of coherent states $\{|\psi\rangle\}$. Actually, we are interested only in the relation between the expansion for $|I\rangle$ and that for the groundstate $|0\rangle$. In the homogeneous case this relation is trivial because of the Galilean invariance: If

$$|0\rangle = \int D\psi Q[\psi] |\psi\rangle, \quad (15)$$

then

$$|I\rangle = \int D\psi Q[\psi] |e^{i\frac{2\pi Ix}{L}}\psi\rangle. \quad (16)$$

In the general case one may think of the state $|I\rangle$ as being obtained from the homogeneous one (16) by semiadiabatic turning on the external potential. The word "semiadiabatic" means that the turning-on time is much greater than the inverse interlevel separation $\propto L$, but is much less than Δ_I^{-1} . Integrating over all the variables except for the long-range phase field Φ we have

$$\langle -I | e^{-H\beta} | I \rangle = A \int D\Phi e^{-S[\Phi]}, \quad (17)$$

where A is some independent of L and I normalization constant, the field $\Phi(x, \tau)$ is determined at $0 \leq \tau \leq \beta$ and satisfies the boundary conditions

$$\int_0^L dx \frac{\partial}{\partial x} \Phi(x, 0) = 2\pi I, \quad \int_0^L dx \frac{\partial}{\partial x} \Phi(x, \beta) = -2\pi I. \quad (18)$$

To the first approximation, semiadiabatic turning on the external potential does not affect Eq.(17), provided β is much greater than the turning-on time.

Calculation of the integral in (17) may be performed by a shift of the variable Φ :

$$\Phi = \Phi_1 + \Phi_0, \quad (19)$$

where the shifting field Φ_0 is chosen to satisfy (below we turn to the variable $y = c\tau$ and correspondingly $y_* = c\beta$)

$$\Delta\Phi_0 = 0 \quad (20)$$

with the boundary conditions (18) ($\Phi \rightarrow \Phi_0$). Then $S[\Phi] = S[\Phi_1] + S[\Phi_0]$ and Φ_1 may be integrated out taking into account that

$$A \int D\Phi_1 e^{-S[\Phi_1]} = \langle 0 | e^{-H\beta} | 0 \rangle = e^{-E_g\beta}. \quad (21)$$

As is easily seen, to meet the boundary conditions Φ_0 should contain a number of vortices (=instantons) with the total charge $2I$. So besides the integration over Φ_1 there should be also an integration over the positions of the instantons: $D\Phi \rightarrow D\Phi_1 \prod_{j=1}^{N_*} dx_j dy_j$, (x_j, y_j) being the position of the vortex j . Generally speaking, there is also summation over the number of instantons, N_* . For our purposes, however, we may confine ourselves to the case $\beta \ll \Delta_I^{-1}$, where the main contribution comes from the N_* corresponding to the minimal possible value of $S[\Phi_0]$. This implies that all vortices are of the minimal possible charge of the same sign. We notice also that in this case $\sin(-i\Delta_I\beta) \approx -i\Delta_I\beta$. Consequently, from (14) and (21) we obtain the relation for Δ_I :

$$\int \prod_{j=1}^{N_*} dx_j dy_j e^{-S[\Phi_0]} \propto e^{-E_I\beta} \Delta_I\beta, \quad \Delta_I\beta \ll 1. \quad (22)$$

Consider the most simple case of the splitting of the two-fold degenerate groundstate level in the system with the gauge phase π , which we will refer to as Aharonov-Bohm (AB) splitting, Δ_{AB} . Upon the elimination of the vector potential by the gauge transformation this case corresponds to $I = \pm 1/2$. It is worth mentioning certain ring systems in which such a case occurs automatically: fermions with the even number of particles; lattice spins with antiferromagnetic interaction and lattice bosons with the positive hopping amplitude, both at the odd number of sites.

For $I = 1/2$ minimal $S[\Phi_0]$ corresponds to Φ_0 with the only one vortex of the charge equal to unity:

$$S[\Phi_0] = i\gamma(x_0) + \frac{\pi^2 \Lambda_s}{2L} \beta + K^{-1} \ln L + o(L^{-1}) , \quad (23)$$

where x_0 is the x -coordinate of the vortex core. Taking into account that

$$\int_0^L dx_0 e^{i\gamma(x_0)} \propto \begin{cases} \text{const} & \text{one defect} \\ \sim \sqrt{L} & \text{disordered system} \\ L & \text{commensurate with integer } \nu \\ 0 & \text{otherwise} \end{cases} \quad (24)$$

from (22) we obtain the estimate for the relative value of the AB splitting.

$$\frac{\Delta_{AB}}{E_{AB}} \propto \begin{cases} 1/L^{1/K-1} & \text{one defect} \\ \sim 1/L^{1/K-3/2} & \text{disordered system} \\ 1/L^{1/K-2} & \text{commensurate with integer } \nu \\ 0 & \text{otherwise} \end{cases} \quad (25)$$

Here $E_{AB} \equiv E_{I=1/2}$. Note that in the disordered case the value of splitting is sensitive to the particular realization of disorder and experiences relative fluctuations of order unity from system to system. Also, in the regular incommensurate case as well as in the commensurate case with a fractional filling factor Δ_{AB} is equal to zero because of the exact cancellation of contributions from instantons with different x_0 's.

For disordered and commensurate systems the laws (25), generally speaking, take place only at sufficiently large L , where K reaches its macroscopic limit. So they need be revised

in the vicinity of the critical points, where renormalization of K is essential and, moreover, is very slow with respect to L . (Such a problem does not emerge in the single-impurity case, since there is no renormalization of K .) To allow for the renormalization we notice that the only place in the effective action (23) where smaller scales of distance may be important is the vortex core energy $E_V(L) = K^{-1} \ln L$, which now is to be estimated more accurately. This may be done just by noting that according to the above-mentioned meaning of the variable w in RG equations it is related to E_V by

$$w^2(L) \propto \begin{cases} L^3 e^{-2E_V(L)} & \text{disordered} \\ L^4 e^{-2E_V(L)} & \text{commensurate} \end{cases} \quad (26)$$

Thus we obtain

$$\frac{\Delta_{AB}}{E_{AB}} \propto w(L) , \quad (27)$$

where $w(L)$ should be taken from RG equations. In particular, at the critical point, where both (8) and (12) yield $w(L) \sim 1/\ln L$, we have

$$\frac{\Delta_{AB}}{E_{AB}} \propto 1/\ln L \quad (\text{at the critical point}) . \quad (28)$$

In the case of commensurate system with a fractional $\nu \propto 1/p$ only the instantons whose charges are multiples of p are relevant. Clearly, this leads to the fact that only some special SC levels are split. Namely, the levels with I 's being the multiples of $p/2$. Consequently, at an odd p split SC levels arise only at the gauge phase π , while at an even p the gauge phase π renders all SC levels degenerate. Estimation of the splitting of the first split SC level ($I = \pm p/2$) is done in the complete analogy to the estimation of Δ_{AB} . The only difference is that the charge of the instanton now is p rather than unity. So we readily get

$$\frac{\Delta_{p/2}}{E_{p/2}} \propto 1/L^{p^2/K-2} \quad \text{commensurate with } \nu \propto 1/p . \quad (29)$$

This result is a generalization of Eq.(25) for the commensurate case. Obviously, Eqs.(27,28) (with $AB \rightarrow p/2$) are also valid.

If we try to estimate by the same technique the splittings of higher SC levels, when more than one instanton is necessary, we have to deal in (22) with a divergency of the integration with respect to y_j 's. This is a standard problem one faces when considers in imaginary time the tunneling between the excited states rather than between ground-state ones. Instead of thinking of a trick of circumventing this difficulty, we prefer to turn to another language of description. In the next section we introduce the effective Hamiltonian, whose relation to the effective action (2) is clearly traceable, and which allows to treat the problem of the SC splitting as well as the hybridization with phonons most naturally.

III. EFFECTIVE-HAMILTONIAN APPROACH

In this section we demonstrate how one may construct an effective Hamiltonian for the SC dynamics and calculate the ratios Δ_I/Δ_{AB} as well as the resonances with the phonons.

In the Fourier-transformed fields the Hamiltonian of the superfluid phase can be written as

$$H_0 = \sum_q \omega_q b_q^\dagger b_q + \frac{\pi}{2L} (v_N \delta N^2 + 4v_I I^2), \quad (30)$$

where $v_I = \pi\Lambda_s$, $v_N = (\pi\kappa)^{-1}$, $b_q^\dagger$ is the standard Bose creation operator, $\omega_q \approx c |q|$ in the long-wavelength limit. For completeness, we have introduced the term associated with the energy change when we add particles to the system, $\delta N = N - N_0$.

We start by noting that the instanton is nothing else but an effective vertex connecting nearest SC. Thus in the most general form we may present the instanton at the point x as

$$H_{int}(x) = \sum_I \left(\frac{V_\Lambda}{L} e^{i\gamma(x)} \hat{O}(b) c_{I+1}^\dagger c_I + h.c. \right), \quad (31)$$

Here $c_{I+1}^\dagger$ is the raising operator of the quantum number I , $\gamma(x)$ is given by Eq.(5), and $\hat{O}$ is an operator acting on the phonon states. The dependence on the coordinate comes from the fact that an intrinsic momentum of the current state in the homogeneous system equals to $I2\pi n$ and the Hamiltonian (31) changes I by 1. Indeed, the SC is obtained by shifting all the

particles in the momentum space by I quanta with the total momentum change $NI \ 2\pi/L$. In disordered systems the particle density is no longer homogeneous and we must replace the average particle density with the local one $n \rightarrow n_0(x)$ according to (5). To satisfy the RG equation for the effective vertex w (see also below) the scaling dimension of $\hat{O}$ should equal to $1 - 1/K$.

As shown by Haldane [2] (and will be demonstrated explicitly below for the particular case of electron backscattering from the impurity) the bosonic field responsible for the current relaxation is

$$\theta(x) = (\theta_0 + \gamma(x)) + i \frac{1}{\sqrt{K}} \sum_{q \neq 0} \sqrt{\frac{2\pi}{|q|L}} \text{sign}(q) (b_q + b_{-q}^\dagger) e^{-iqx}. \quad (32)$$

and the original particle field is given in terms of exponential functions of $\theta(x)$. By noting that the phase θ_0 is the canonical momentum for I , the resulting effective Hamiltonian may be rewritten as

$$H_{int}(x) = \sum_{-\pi < \theta_0 \leq \pi} \left(\frac{V_\Lambda}{L} e^{i(\theta_0 + \gamma(x))} \hat{O}(b) c_{\theta_0}^\dagger c_{\theta_0} + h.c. \right), \quad (33)$$

Clearly, the first two terms in the exponent are those in the definition of $\theta(x)$ given above, and we identify the operator part of the vertex $\hat{O}$ in Eq.(31) with the $q \neq 0$ components of the phase field $\theta(x)$

$$\hat{O}(b) = \frac{L}{\Lambda} \exp\{i\theta_{q \neq 0}^{(\Lambda)}(x)\}, \quad (34)$$

where the first factor is necessary for the correct scaling of $\hat{O}$. The sum over momentum has an upper cutoff at some scale $q \leq 2\pi/\Lambda$ where one has to define the "bare" amplitude V_Λ which depends on the microscopic parameters of the initial Hamiltonian and the cutoff. One can not obtain the exact value of the parameter V_Λ from this analysis (except for the scaling behaviour mentioned above). The other way of deriving the effective Hamiltonian mathematically rigorously is by mapping the effective Lagrangian Eqs.(2,4) to the equivalent sine-Gordon model [4], which is in fact done by Eq.(33) since $H_{int} \sim \cos \theta(x)$.

There is yet another simple microscopic derivation of Eq.(34) for the particular case of electron backscattering from the impurity potential. In the bosonic representation the

original right- and left-moving electron fields (labeled by the index $p = \pm 1$) have the form [9]

$$\Psi_p^\dagger(x) = \frac{1}{\sqrt{\Lambda}} \mathbf{U}_p e^{-ipk_F x} \exp \left\{ \sum_q \sqrt{\frac{2\pi}{|q|L}} \text{sign}(q) C_{pq} (b_q + b_{-q}^\dagger) e^{-iqx} \right\}, \quad (35)$$

$$C_{pq} = \begin{cases} \cosh(K), & pq > 0 \\ \sinh(K), & pq < 0 \end{cases}$$

where the Fermi momentum k_F equals to $\pi N/L$, and $\mathbf{U}_p$ is the ladder operator increasing the number of p -particles by one. If we substitute these expressions into the backscattering Hamiltonian, $H_B = V_B \Psi_r^\dagger \Psi_l + h.c.$, which is responsible for the current relaxation on impurities, we immediately arrive at the effective Hamiltonian Eq.(33,34) [13]. An important new point here is that it is valid for any 1D system in the superfluid phase, provided one uses the proper definition of the quantum number I , which has the particular meaning of the difference between the number of left- and right-moving particles for fermions only.

A short physical discussion is in order here. One may consider the effective Hamiltonian (31) as resulting from the coupling between the SC states and sound waves of the form

$$\sim c_I^\dagger c_I \frac{Ic}{\sqrt{K}} \sum_{q \neq 0} \sqrt{\frac{2\pi |q|}{L}} \text{sign}(q) (b_q - b_{-q}^\dagger) e^{-iqx}. \quad (36)$$

As is well known, this interaction can be eliminated by the standard on-site polaronic transformation in expense of the additional operator structure $\hat{O}$ in the hopping term. Thus any SC state causes a long-range deformation in the phonon field, and this deformation is growing linearly with the quantum number I . In the fermion system this is equivalent to "dressing" each extra left/right-moving particle by density fluctuations. Now any attempt to change the SC state will result in the readjustment of the phonon vacuum to the new value of I . It turns out that the dominant contribution to the effective low-energy vertex is due to this reconstruction of density fluctuations rather than due to the "bare" process of current relaxation which defines the high-energy prefactor here. As K goes to zero the coupling between the SC state and density fluctuations diverges. This limit describes the case

of weakly interacting bosons and quasi 1D superconducting rings (see section V), where even the lowest SC state (which still has a finite momentum!) involves *large* number of particles with *small* momenta moving in the same direction. Now the process of current relaxation is through the backscattering of many particles simultaneously, rendering the low-energy vertex extremely small. Thus a one-particle description of SC in terms of the number of left- and right-moving carriers is clearly misleading for $K \rightarrow 0$; the relevant language in all cases being that based on the quantum number I .

One may now straightforwardly check that the renormalization group equations in the disordered case are correctly described by the effective Hamiltonian. The RG equation for the vertex is obtained by a simple averaging of the operator $\hat{O}$ over the momentum range $2\pi/\Lambda' < |q| < 2\pi/\Lambda$ where $\Lambda' = \Lambda + d\Lambda$. In the second order perturbation theory the renormalization of the superfluid energy is due to the virtual transition of the system to the neighbouring current state with the simultaneous emission of the sound mode with $2\pi/|q| \in (\Lambda, \Lambda')$

$$\delta E_I = -(V/L)^2 \frac{1}{K} 2 \int_{(\Lambda, \Lambda')} \frac{dq}{q} \left[\frac{1}{2\pi c/\Lambda + E_{I+1} - E_I} + \frac{1}{2\pi c/\Lambda + E_{I-1} - E_I} \right]. \quad (37)$$

where $E_I = 2\pi v_I I^2/L$. By expanding the denominators up to the second order in $1/L$ we find the RG equation for the current velocity:

$$\frac{d \ln v_I}{d \ln \Lambda} = -\frac{4V^2 \Lambda}{\pi^2 c^2 K^2 L}. \quad (38)$$

The renormalization of the sound velocity is due to the virtual transitions from I to $I \pm 1$ with the simultaneous absorption of the sound wave and emission of the high-frequency mode. In full analogy with the previous calculation we have

$$\frac{d \ln c}{d \ln \Lambda} = -\frac{2V^2 \Lambda}{\pi^2 c^2 K^2 L}. \quad (39)$$

As expected, one may neglect the renormalization of v_I and c in the superfluid phase when dealing with the single impurity scattering (the r.h.s. of Eqs.(38,39) is $1/L$ in this case).

For disordered potential the effective Hamiltonian is an obvious sum over the impurity positions, and in Eq.(31) one has to replace $V \rightarrow \sum_{x_i} V e^{i\gamma(x_i)} \sim V(L/\Lambda)^{1/2}$. Now the

scaling equation for the vertex is changed to $dV/d\ln\Lambda = V(3/2 - 1/K)$, in agreement with Eq.(12). Correspondingly, in Eqs.(38,39) one has to replace V^2 by V^2L/Λ . By introducing the dimensionless coupling parameter $w = 2V/\pi cK$ we rewrite the RG equations as

$$\begin{aligned}\frac{d\ln v_I}{d\ln\Lambda} &= -w^2, \\ \frac{d\ln c}{d\ln\Lambda} &= -w^2/2,\end{aligned}\tag{40}$$

or, using the universal relations $c = v_I K = v_N/K$,

$$\frac{dK}{d\ln\Lambda} = w^2 K; \quad v_N = \text{const}.\tag{41}$$

Thus we have reproduced the RG equation (12) of section II.

In exactly the same manner one can construct an effective Hamiltonian for the commensurate system. Let $n = 1/p$. Then the vertex connecting the states I and $I \pm p$ is defined by

$$H_{comm} = \sum_I \int \frac{dx}{\Lambda} \left(\frac{V}{\Lambda} e^{ip\theta_{q \neq 0}(x)} c_{I+p}^\dagger c_I + h.c. \right).\tag{42}$$

Due to the translational symmetry of the commensurate problem there is no renormalization of the sound velocity. For the same reason only states with zero total momentum are hybridized with the topological excitations and e.g. the correction to the superfluid energy is defined by virtual transitions to the state with two high-frequency modes $q_1 = -q_2 \in (\Lambda, \Lambda')$. Now

$$\frac{dK}{d\ln\Lambda} = p^6/16 w^2; \quad c = \text{const}.\tag{43}$$

The coefficient in front of the w^2 term can be absorbed into the redefinition of w (cf. Eq.(8)).

The effective-Hamiltonian approach reduces the discussion of the superfluid - insulator transition to that of a "particle" localization at a given site I . In the ground state (when $\Lambda = L$) the current Hamiltonian is identical to the tight-binding model with the hopping amplitude V_L/L in the static external potential given by the supercurrent energy

$$H(L) \approx \sum_I \frac{2\pi v_I}{L} (I - \phi/\phi_0)^2 c_I^\dagger c_I + \sum_I \left(\frac{V_L}{L} c_{I+1}^\dagger c_I + h.c. \right).\tag{44}$$

If under scaling one finds $V_L \ll v_I$, then the potential energy prevents the current state from delocalization, and we have perfect metallic conductance. Only at some special values of the magnetic flux through the ring (e.g. integer or half-integer flux), when the states I and $-I$ are in resonance, the current is delocalized between the two states, but the energy splitting of the resonance is much less than $2\pi v_I/L$.

The superfluid-insulator transition is described as a delocalization of the current state, when $V_L \gg v_I$. In this regime the topological invariant is delocalized in a band and I is no longer a well-defined quantum number [13].

Let us now calculate the splittings between the resonating states. If we neglect the hybridization with phonons and work within the Hamiltonian (44), then the splitting is given by the perturbation theory of $2I$ -th order in V_L

$$\begin{aligned}\Delta_I &= 2 \frac{V_L^{2I}}{L} \prod_{l=1-I}^{I-1} \frac{1}{2\pi v_I(I^2 - l^2)} \\ &= \frac{4\pi v_I}{L} \left(\frac{V_L}{2\pi v_I} \right)^{2I} \frac{1}{[(2I-1)!]^2} .\end{aligned}\tag{45}$$

If the ring is in the flux $\phi = \phi_0/2$ then the splitting between the states I and $1-I$ is described by the same expression (45) with the replacement $I \rightarrow I - 1/2$.

In deriving Eq.(45) we considered only the path from I to $-I$ which corresponds to the ground state of the bosonic field in *all* intermediate states. In the general case one has to consider the possibility of exciting/absorbing an arbitrary number of sound waves in any intermediate state. To this end we slightly "undress" the ground-state vertex and write the final low-energy Hamiltonian in the form

$$H_{eff} = \sum_I \frac{2\pi v_I}{L} (I - \phi/\phi_0)^2 c_I^\dagger c_I + \sum_I \left(\frac{V_L}{L} \frac{e^{i\theta}}{\langle e^{i\theta} \rangle} c_{I+1}^\dagger c_I + h.c. \right) .\tag{46}$$

which is cutoff independent. First, let us consider the case of single impurity, when the spatial dependence of V_L is $\delta(x - x_i)$ at the impurity position x_i . Using standard properties of bosonic operators we readily calculate all the matrix elements

$$\langle \{N_q\} | \frac{e^{i\theta}}{\langle e^{i\theta} \rangle} | \{N_q + S_q\} \rangle = \prod_q r(N_q, N_q + S_q, \frac{2\pi}{LK |q|}) ,$$

$$r(N, N+S, x) = \frac{e^x x^{S/2}}{\sqrt{N!(N+S)!}} \sum_{n=0}^{\infty} \frac{(-1)^n (N+S+n)!}{x^n n!(n+S)!} \begin{cases} 1 & S \geq 0 \\ (-1)^S & S < 0 \end{cases}. \quad (47)$$

In principle, the problem of the low-energy spectrum is solved now, because we know explicitly all the matrix elements. In a large system the coupling V_L is macroscopically small, and apart from extremely narrow resonances at $1/K = \text{integer}$, when one has to solve a finite-size secular equation, it is sufficient to restrict ourselves to the perturbation theory in V_L . Below we calculate analytically the energy splitting of the first SC state.

First, we note that the coupling to phonons does not modify the splitting of the ground state in a ring with $\phi = \phi_0/2$, simply because there are no intermediate states in this case (see Eq.(25))

$$\Delta_{AB} = \frac{4\pi v_I}{L} (V_L/2\pi v_I). \quad (48)$$

With the coupling to phonons taken into account the expression for the first SC splitting acquires the form

$$\Delta_1 = \Delta_{AB}^2 \frac{L}{4\pi v_I} \sum_{\{q, S_q\}} \frac{\Pi_q (-1)^{S_q} |r(0, S_q, \frac{2\pi}{LK|q|})|^2}{1 - \sum_q \frac{LK|q|}{2\pi} S_q}. \quad (49)$$

Next we introduce the integer variable M and write down the spectral representation for the δ -symbol $\delta(M - \sum_q (L | q | / 2\pi) S_q)$. This allows to calculate the contribution of each phonon mode separately

$$\begin{aligned} R &= \frac{4\pi v_I \Delta_1}{L \Delta_{AB}^2} = \int_{-\pi}^{\pi} \frac{dt}{2\pi} \sum_{M=0}^{\infty} \frac{e^{-iMt}}{1 - KM} \exp \left\{ -\frac{2}{K} \sum_{p=1}^{\infty} \frac{e^{ipt}}{p} \right\} \\ &= \sum_{M=0}^{\infty} \int_{-\pi}^{\pi} \frac{dt}{2\pi} \frac{e^{-iMt} (1 - e^{it})^{2/K}}{1 - KM}. \end{aligned} \quad (50)$$

Straightforward evaluation of this expression gives the final answer

$$R = \frac{\pi/K}{\sin(\pi/K)} \frac{\Gamma(1 + 2/K)}{\Gamma^2(1 + 1/K)}. \quad (51)$$

Note that $R = 1$ if we neglect the hybridization with phonons.

Thus we found that for $1/K = \text{integer}$ the splitting diverges due to the resonance between the SC and phonons. For small V_L this divergence is accurately described by Eqs.(50,51)

except for a tiny region of order V_L/v_I around integer values of $1/K$. However in this region the ratio R is already very large, and the dominant contribution comes from the coupling to a small group of phonon states with the same energy $\approx 2\pi v_I/L$. For the particular case of $1/K \approx 2$, which will be compared in the next section with the exact numerical diagonalization, one has to take into account the states $|N_q = 2\rangle$, $|N_{-q} = 2\rangle$, $|N_q = 1, N_{-q} = 1\rangle$, $|N_{2q} = 1\rangle$, and $|N_{-2q} = 2\rangle$ where $q = 2\pi/L$. The hybridization matrix elements between these states and SC squared are 2, 2, 4, 1, and 1, correspondingly. By writing a secular equation for this group of levels we obtain a complete picture of level crossings between the SC and sound modes.

Quite generally, at the point of resonance *all* energy splittings are of the same order $\Delta_I \sim \Delta_{AB}$. Indeed, for integer $1/K$ the instanton is always between the states with the same energy (for the combined system SC + phonons), so the splitting is given by the amplitude of a single transition Δ_{AB} no matter how many instantons are required to connect the resonating SC states.

This technique may be applied for calculating the energy splittings in commensurate systems as well. An important difference with the impurity case is that now the supercurrent energy change is $\delta E_{I-p,I} = 2\pi v_I(2pI - p^2)/L$ and resonating states are given by $I = p/2 \pm mp$, and only sound waves with the zero total momentum are hybridized with the SC. We find that for commensurate systems we always have a resonance between the SC and sound at the point of superfluid-insulator transition. Indeed, for $K = p^2/2$ the energy change $\delta E_{I \pm p,I} = 2\pi c/L \cdot 2(2n - 1)$ is given by even number of the phonon energy quanta.

Strictly speaking, there is only one cutoff-dependent parameter V_L (apart from the Luttinger Liquid parameters c and K) which controls the low-energy spectrum of the system. Once this parameter is fixed, say from the Δ_{AB} splitting, all higher energy splittings and level crossings can be found explicitly. However in the disordered system we face a problem. By construction the low-energy vertex $V_L(x)$ in this case is a random function of coordinate, and depends on the particular disorder realization in the sample. Now, even if we define the mean value of $\langle V_L \rangle$ from the lowest splitting Δ_{AB} , we still are uncertain about the inte-

grals $\int_0^L dx e^{\pm i 2 \pi m x / L} V_L(x)$ which define the hybridization matrix elements with the phonon modes and take care about momentum non-conservation in disordered system. The only thing which is certain is that for the lowest levels with $m \sim O(1)$ these integrals have random values of order $\langle V_L \rangle$ in magnitude. It makes the direct comparison between the theoretical calculation (51) and the numerical result for R inconclusive, unless additional parameters are introduced into the theory to account for the random variations of the above-mentioned integrals (see more in the discussion of the numerical results in section IV).

The above discussion dealt with the properties of an isolated 1D ring. Obviously, by considering the lowest energy levels we lack any kind of dissipation in the system. In the ideal picture the current state I obtained, e.g., by nonadiabatic switching on the magnetic flux through the ring will persist forever, or demonstrate undamped oscillations with the frequency Δ_I . The most effective dissipation mechanism is due to the coupling between the ring and substrate. One has to take an insulating substrate in order to eliminate the conduction electrons which form a thermal bath with the largest low-frequency density of states, and thus are the most dangerous source of decoherence. To couple the SC states to phonons of the substrate we have to consider the variation of H_{eff} originating from the lattice long wave-length distortion $u(x)$. The effect of $u(x)$ is two-fold: to change the kinetic and potential energy. For example, the distortion with the wave-length of order L (which can absorb the energy difference $\delta E_{I,I-p}$) may influence the total length of the ring $\delta L \sim Lu$ and result in the u -dependent terms both in the kinetic and potential energy. For disordered sample the positions of impurities $\{x_j\}$ depend on the substrate deformation giving $\delta V_L \sim V_L u$. One may also take into account the dependence of K (or the particle-particle interaction) on the inter-particle separation.

It is easy to check that both the nondiagonal and diagonal couplings give the same value of the relaxation time τ . We write the diagonal interaction Hamiltonian as

$$H_{diss} \sim \alpha \frac{v_I}{L} u \sim \alpha \frac{v_I}{L} \left[\sum_k \sqrt{k/Ms} (d_k + d_k^\dagger) \right], \quad (52)$$

where $d_k^\dagger$ are the bulk phonon creation operators, α is the dimensionless parameter depending

on the microscopic details and geometry of the particular system, M is the substrate ion mass, and s is the bulk sound velocity. Now, using standard results for the relaxation time in a double-well system coupled to phonons (see e.g. [16]) we find

$$\tau^{-1} \sim \left(\frac{V_L \alpha v_I}{L^2} \right)^2 \frac{|E_{I,I-p}|}{\rho s^5} \sim \frac{V_L^2 \alpha^2 v_I^3}{\rho s^5 L^5}, \quad (T = 0). \quad (53)$$

Here ρ is the substrate density. If we couple the substrate to the nondiagonal terms in Eq.(46) by expanding

$$V_L/L \rightarrow V_L/L (1 + \alpha' u),$$

then the final result for τ^{-1} coincides with the Eq(53).

The relaxation rate should be much less than the level splitting Δ_I if we want to observe the coherent oscillations. In the mesoscopic limit we find $\Delta_I \tau \sim V_L^{2(I-1)} L^4$. Substituting here the scaling behaviour of V_L , we obtain the criteria defining which levels will demonstrate coherence at $T = 0$

$$I < \begin{cases} (1+K)/(1-K) & \text{single impurity} \\ (2+K)/(2-3K) & \text{disordered} \\ p^2/(p^2-2K) & \text{commensurate} \end{cases}. \quad (54)$$

We see that the broadening of the two lowest doublets Δ_{AB} and Δ_1 is always small. Very close to the superfluid-insulator transition we can safely neglect dissipation for all levels. To get a better feeling of how long the relaxation time could be in a real system, we estimate τ^{-1} from Eq.(53) using $\gamma \sim 1$, $v_I \sim s$, $L \sim 10^4 A^o$, and the maximum possible $V_L \sim v_I$. For this particular choice of parameters $\tau \sim \rho L^5 \sim 1$ week.

IV. NUMERICAL SPECTROSCOPY

In this section we report exact numerical spectroscopic results for 1D finite-size boson Hubbard models. The Hamiltonians we examine can be written in the most general form as (all the parameters are scaled in the units of the hopping amplitude)

$$H = \sum_{i=1}^{N_a} \left\{ -(a_i^\dagger a_{i+1} + \text{h.c.}) + \epsilon_i n_i + \frac{U}{2} n_i (n_i - 1) + V n_i n_{i+1} \right\}. \quad (55)$$

Here a_i is the annihilation operator on the site i ; $n_i = a_i^\dagger a_i$, and $\{\epsilon_i\}$ are random numbers uniformly distributed over the range from $-W/2$ to $W/2$. (In the single-impurity case all the ϵ_i 's are equal to zero except for ϵ_1 .) The ring topology implies that the sum of the site indices is understood modulo N_a , the total number of sites. The gauge phase π , if necessary, is introduced by changing the sign of one of the hopping amplitudes (anti-periodic boundary condition).

For the calculation of the scalings of SC splittings with increasing N_a we use hard-core models ($U = \infty$) which at a given N_a have much less Hilbert-space dimension as compared to the full models ($U \neq \infty$), and thus allow consideration of larger systems. In this case the term with V is necessary to control the parameter K . In the case of a full model, where this term does not introduce any qualitative difference with respect to the on-site term U , we set $V = 0$.

Our diagonalization procedure [22] is arranged as follows. First, starting from some trial wavefunction we calculate a certain number of the lowest energy levels and corresponding wavefunctions by modified Lanczos method [19]. The set of approximate eigenfunctions is reconstructed from Relay's tridiagonal matrix [20], and the trial wavefunction is expanded in it. As is known, the set inevitably involves a substantial number of spurious states, due to numerical errors. These states, however, may be easily identified by their negligible contribution to the expansion of the trial wavefunction. Upon exclusion of the spurious states the set is subjected to the orthogonalization and correction by Newton method. The relative (with respect to a characteristic interlevel spacing) errors of the energy level calculation are typically of order $10^{-13} \div 10^{-11}$ for the groundstate, and of order $10^{-9} \div 10^{-5}$ for some ten first excited states. The problem of degeneracy is solved by repeating the procedure with a new trial function, chosen to be orthogonal to all previously obtained eigenstates. The combined set of eigenfunctions is again orthogonalized and corrected by Newton method.

In the case of the regular system we take advantage of the translational symmetry to

proceed separately for each momentum sector.

Now we turn to the results. To get an idea of how the spectra look like in this or that case, we show the pictures of spectrum evolution with increasing on-site disorder (Fig. 1) and on-site interaction (at a fixed disorder) (Fig. 2), and as a function of the nearest-neighbor interaction in the case of commensurate hard-core model (Fig. 3). All the energies are reckoned from the ground states. Besides, in the case of disordered system we found it convenient to normalize the energy values in such a way that the middle of the first phonon doublet always corresponds to unity. With this normalization the value of the first SC coincides with K^{-1} . The level identification is performed in accordance with [22], where the numerical spectra for a disordered 1D ring were reported for the first time.

Though it is not directly related to the subject of the present paper, it is interesting to discuss the behavior of the hard-core model at large negative V 's. From general considerations it is clear that at a certain negative $V = V_*$ there should occur an ordinary condensation into a macroscopic drop. So that at $V < V_*$ the first $N_a - 1$ energy levels (associated with the motion of the drop as a whole with momenta $= 1, 2, 3, \dots, (N_a - 1)$) should lie very close to the ground-state one (corresponding to the zero-momentum motion of the drop). It is precisely what is seen in Fig.3. From the symmetry of the hard-core model (isomorphism with the spin-1/2 system) it immediately follows that in our case $V_* = -2$ is the point where, speaking the spin language (isotropic ferromagnetic Heisenberg Hamiltonian), the transition to the ferromagnetic ground state takes place. To minimize the energy below this point a system with fixed z -projection of the full spin (fixed number of particles in our case) would desire to be divided into two parts: with spins up and spins down, respectively (a part with empty sites and a part with completely occupied ones). In a regular system this symmetry breaking is hidden by translation invariance, but may be revealed by calculating some relevant correlator. Consider e.g. $Q_3 = \langle n_{i_1} n_{i_2} n_{i_3} \rangle / \langle n_{i_1} \rangle^3$, where $|i_1 - i_2| = |i_2 - i_3| = |i_3 - i_1| = N_a/3$. In the macroscopic limit we have $Q_3 = 1$ for homogeneous liquid and $Q_3 \rightarrow 0$ for a condensed one. Our calculations for $N_a = 18$ with the filling factor $\nu = 1/2$ yield $Q_3 \approx 0.90$ at $V = -1.8$ and $Q_3 \approx 0.14$ at $V = -2.3$.

In connection with the transition at $V = -2$ in the hard-core model it is worth mentioning that just at this point there exists an analytical Bethe-ansatz treatment recently reported by Sutherland [21], which reveals the appearance of macroscopic bound complexes.

The scaling of the single-impurity AB-splitting (Fig. 4) is in excellent agreement with the macroscopic theory. Here and below the dashed line ("theory") is plotted in such a way that it passes through the last (maximum N_a) point and has a slope corresponding to Eqs.(25, 29), the value of K being obtained from the numerical spectrum at this point (sound velocity can be taken from the first phonon level while Λ_s is available e.g. from the first SC).

The analogous results for the disordered system (Fig. 5) demonstrate a good agreement with the theory only at larger N_a . In fact, this is what one might expect to get bearing in mind that the value of K we deal with (~ 0.5) is not very far from the critical one ($= 2/3$), so that the renormalization of K due to the disorder may be significant. In accordance with Eq.(25) the decrease of Δ_{AB} with increasing N_a at $K \sim 0.5$ is rather slow. Hence, to reveal it against the statistical dispersion, characteristic of the disordered case, an extensive averaging is required.

Fig. 6 shows the scaling of Δ_1 in the commensurate case. The results at $V = -0.5$ are in excellent agreement with Eq.(29), while at $V = -1.5$ there is a slight deviation (which, however, does not exceed 15%). Since this deviation has a tendency to decrease with increasing N_a , it may be accounted for as a small-size effect.

Fig. 7 shows the scaling of Δ_1 for the commensurate system at the critical point and demonstrates the excellent agreement with the prediction of the macroscopic theory that the plot will approach a straight line as $N_a \rightarrow \infty$. It should be noted that for the half-filled 1D hard-core model the critical point (corresponding to $V = 2$ where the model is isomorphic to the isotropic Heisenberg antiferromagnetic spin-1/2 chain) was extensively studied [23]. In particular, it was shown that some excited levels demonstrate logarithmic ($\sim 1/\ln N_a$) relative shifts of their positions as compared to the macroscopic limit. Speaking superfluid language [25], we understand that these are the states that have either non-zero

topological quantum number I , or extra particles/holes. (Purely phonon excitations can not demonstrate such shifts since the sound velocity is irrenormalizable quantity.)

To verify the correctness of the effective-Hamiltonian approach Eqs.(31,34) we calculate numerically the relation between the SC splittings Δ_1 and Δ_{AB} for the disordered ($W = 0.25$) system of 7 bosons on 11 sites. The coupling parameter U varied in the range $2.6 \leq U \leq 6.2$ where the first SC level crosses the group of phonon levels with energies $\approx 4\pi c/L$. The numerical ratio $R = 2E_1\Delta_1/\Delta_{AB}$ is shown in Fig.8. Unfortunately, the direct comparison with the theoretical expression (51) is not very conclusive here because it was calculated by assuming that *all* 5 phonon levels with energies $\approx 4\pi c/L$ are degenerate. However in a very small ring this is not the case even approximately as is clearly seen in Fig.2. The single resonance is split and in the parameter range $U < 3$ other multi-phonon states (also strongly split) practically wash out any sign of the phonon spectrum quantization in multiples of $2\pi c/L$. Under these conditions one may not expect to see any reasonable agreement between the calculated ratio Eq.(51) and the numerical results on such small systems.

One may try to improve the theory by adjusting the calculated ratio (51) to the actual positions of the phonon levels "by hand", e.g., by applying the procedure

$$R \rightarrow R - \sum_{\{q, S_q\}}^{group} \prod_q (-1)^{S_q} \left| r(0, S_q, \frac{2\pi}{LK | q |}) \right|^2 \left(\frac{1}{1 - KM(\{q, S_q\})} - \frac{E_1}{E_1 - E_{ph}(\{q, S_q\})} \right), \quad (56)$$

where the sum is over any group of levels characterized by $\{q, S_q\}$ with the matrix elements given by (47) (more precisely the ratio has to be calculated from the secular equation for the same group of levels). This looks like a reasonable approach to the single-impurity case. Still, in the disordered case we have further uncertainty in the values of the matrix elements because of the integrals $\int dx V_L(x) \exp\{\pm i\gamma(x) \pm i2\pi mx/L\}$, which depend on the particular disorder realization. Obviously, a single value of $\int dx V_L(x) \exp\{i\gamma(x)\}$ taken from the amplitude Δ_{AB} is not sufficient to fix all these integrals for different phonon modes. In the general case we have to treat these integrals as random values, which for the lowest modes

are of order $\langle V_L \rangle$ in magnitude. The avoided level crossing in Fig.2 tells unambiguously that all the symmetries are violated, and SC is coupled to the left- and right-moving phonons with different matrix elements. On the other hand we see clearly in Fig.8 that the splitting Δ_1 has extended resonance structure and the ratio R is typically much larger than unity in qualitative agreement with the theoretical prediction (51).

Finally, we touch upon a problem of an extrapolation of the finite-size exact-diagonalization results to the larger system sizes. In 1D this problem is especially important since the finite-size cross-over regions in the vicinities of the critical points have only logarithmical macroscopic smallness. (For a typical example see the spectrum in Fig. 3 demonstrating quite a smooth behavior at $V = 2$.) Obviously, the extrapolation can be performed with the RG equations. As the parameter K naturally follows from the spectrum, the only problem is to obtain y . This can be done by fitting the behavior of K as a function of N_a with RG equations. The details and results of such an approach see in [24].

V. POSSIBLE REALIZATIONS OF 1D SUPERCURRENT STATES

In this section we discuss in some detail different systems and experimental setups which allow to study SC states in a ring geometry. The most obvious choice is a real 1D electron wire of mesoscopic size. Recent experimental techniques [26] produce rings with $L \sim 10^4 A^o$ and only few transverse levels (in ideal case one) below the chemical potential μ . The experiments can be done on ensembles as well as on single rings [27]. In a static experiment one may study SC by measuring the so-called persistent current, which is defined as

$$j = -\frac{1}{2\pi} \partial F / \partial \varphi \quad (\varphi = \phi / \phi_0) , \quad (57)$$

where F is the free energy. At zero temperature $j = 2v_I/L(I - \varphi)$, where I minimizes the potential energy $E_I(\varphi)$. In ideal system $j(\varphi)$ is a saw-like function with abrupt change of the current from $-v_I/L$ to v_I/L at $\varphi = m + 1/2$, when the two energy levels $I = 0$ and $I = 1$ are degenerate. The singularity is changed to crossover when we incorporate the finite

energy splitting Δ_{AB} into the problem. A trivial solution of the two-state Hamiltonian gives now

$$j(\varphi) = \frac{2v_I}{L} \delta\varphi \left(\frac{2\pi v_I/L}{\sqrt{(4\pi v_I \delta\varphi/L)^2 + \Delta_{AB}^2}} - 1 \right) ; \quad \delta\varphi = \varphi - 1/2 . \quad (58)$$

We see that the crossover region is very narrow $|\delta\varphi| \sim L\Delta_{AB}/2v_I$ in the superfluid phase. Clearly, such a ring may work as a very sensitive magnetometer capable of measuring the flux with the accuracy $\sim \phi_0(v_I/L\Delta_{AB}) \ll \phi_0$. (An interesting experimental setup was proposed recently in [28], where the ring conductance in the flux is probed by two weak tunnel junctions at a distance $L/2$. In this case the Josephson critical current at zero voltage changes dramatically at $\varphi = 1/2$.)

The static picture just described is valid provided the time of experiment is much longer than the relaxation time τ (see Eq.(53)), or if the flux was switched on adiabatically $d \ln \varphi / dt \ll \Delta_{AB}^{-1}$. For large enough systems the both requirements might be violated. By inserting nonadiabatically the flux $\varphi \approx m$ (same arguments go through for $\varphi = 1/2$ -integer) we prepare the SC state $|I = m\rangle$. Then, according to the standard two-state dynamics, the current starts oscillating with the frequency $[(8\pi n v_I/L)^2 + \Delta_I^2]^{1/2}$. In a more sophisticated experiment one may study the echo signal from the ring. Actually, the echo or the burned hole experiment on the ensemble of rings with different sizes seems to be the most appropriate one, because the wide spread in Δ_I and φ for individual rings makes it difficult to see the resonant absorption of the electro-magnetic field. We are not going into details here because this case is identical (with clear modifications for the distribution function) to that of the two-level systems in amorphous materials (see e.g. the review article [29]).

The other system described as a 1D Luttinger Liquid at very low temperatures is in fact a 2D electron gas in the FQHE state. In the FQHE state all bulk excitations are frozen at temperatures much less than the energy gap in the collective spectrum of interacting electrons, and the only low-energy degrees of freedom left in the incompressible electron liquid are the edge excitations which form 1D chiral Luttinger Liquid [30–33]. Here we will

discuss the sample with the Carbino disk geometry where the right- and left-moving fields are described by the inner and outer edge currents correspondingly, and the relaxation process is via quasiparticle tunneling between the edges through the bulk at the point of constriction (or many constrictions if we are interested in the disordered case). The crucial parameter K^{-1} in this case equals to the filling factor of the FQHE state ν [30], and for $\nu = 1/(2k+1)$ (when there is only one channel on each edge) we are in the insulator phase, that is in the *macroscopic* sample $L \rightarrow \infty$ the constriction is equivalent to cutting the ring. One may argue that our results do not apply to this case because we concentrated on the superfluid phase only. However in all our calculations done in sections II, III the only assumption used was that the dimensionless vertex w is small. In the superfluid phase this was true even for a large microscopic "bare" vertex. Suppose now that we have a *finite-size* ring with extremely small "bare" value of w , so that even for large K the scaling toward larger values of w still leaves $w(L)$ small, and we actually never enter the true insulating phase. In this case all our results remain intact and directly apply to the experiment. Moreover, large values of K might be an advantage because we need a ring with the distance between the edges d much large than the magnetic length l_H , in order to be able to speak about well-defined edge excitations separated by inert bulk. The tunneling rate w goes to zero exponentially with the distance d , and for not too narrow disks will be unobservable if not for the increase of w under the scaling when K is large.

The other advantage of the Hall system is that it is not sensitive to small deviations of the magnetic field from special values, and the width of the disk d does not constitute any problem when an external magnetic field is varied to change the flux through the inner edge. Obviously, the above discussion of possible experimental setups for studying the dynamics of supercurrent states is in order here. One can also make use of the spatial separation between the left- and right-moving currents (which is not the case in really 1D wires). In the Hall system current relaxation is connected with the charge transfer between the edges, which allows to follow, e.g., current oscillations, by monitoring the charging effects in the vicinity of the sample.

In all examples discussed so far the parameter K was larger than unity (in 1D conducting wires due to Coulomb repulsion between the electrons we expect $K > 1$). Consider now a 3D superconducting ring having the two spatial dimensions d_1 and d_2 (defining the sample cross-section) much smaller than the ring circumference L . The question we address here is the size dependence of the supercurrent relaxation/oscillation in such a ring. The naive answer is: since the relaxation is via vortex line tunneling across the constriction (or vortex ring nucleation inside the ring with subsequent expansion to the sample boundaries), then the process will depend on d_1 and d_2 , but not on L because it happens locally right at the point of deliberately introduced constriction and none of the vortex sizes exceeds the cross-section diameter. As we demonstrate below, this answer is absolutely wrong, and the relaxation process depends explicitly on the ring length L .

At temperature $T \ll T_c$ (we assume that the superconducting energy gap is nonzero and of order T_c all over the Fermi surface) we describe the superconductor by effective action for the phase of the order parameter which has the same form as Eq.(2)), except that now the integration over dx ought to be understood as $dx_1 dx_2 dx_3$. Let us go to even lower temperatures $T \ll 1/md^2$ (we assume that the cross-section diameter d is larger than the coherence length ξ), when the transverse excitation modes are frozen, and the effective action can be further integrated over dx_1 and dx_2 . What is left is the familiar 1D effective action with $\sqrt{\Lambda_s/\kappa} \sim v_F$ and $\Lambda_s, \kappa \sim (d/a)^2$. Clearly, in this case we deal with extremely small values of K

$$K \sim (a/d)^2 \ll 1, \quad (59)$$

and the effective 1D system is indeed in the superfluid phase. Thus we find that the rate of current oscillations will be proportional to

$$\ln \Delta_{AB}/E_{AB} \sim -(d/a)^2 \ln L/d \quad (60)$$

according to our general arguments which lead to (25) (we used the fact that $1/K \gg 1$).

Till now we completely ignored the effects of Coulomb interaction between the particles, so the result (60) rather applies to the neutral superfluid like He-II in a narrow long tube.

One might also think about the experiment where the superconducting ring is placed on top of the insulator-metal plate, which will screen long-range Coulomb forces. However if the interaction between the charges remains unscreened, the result (60) has to be essentially modified. The spectrum of collective modes in a 3D superconductor in the limit $q \rightarrow 0$ is given by [34] (see also [3])

$$w_q^2 \sim v_F^2 q^2 \Pi(0) V_C(q) ; \quad (61)$$

and for $V_C(q) \sim e^2/q^2$ has a plasmon gap $w_p^2 = ne^2/m \sim v_F^2 \kappa_p^2$, where $\Pi(0)$ is the static polarizability, and κ_p^{-1} is the normal-metal screening radius.

Consider now the limit $L^{-1} \leq q \ll d^{-1}$, which corresponds to our quasi-1D geometry as discussed above. In this momentum range we almost recover back the sound-like energy spectrum for phase fluctuations

$$w_q^2 \sim v_F^2 q^2 (\kappa_p d)^2 \ln qd . \quad (62)$$

Weak logarithmic dependence of the sound velocity and parameter K which is now given as

$$K^{-1} \sim \frac{d}{\kappa_p a^2} \frac{1}{\ln^{1/2}(qd)} \quad (63)$$

can not change the theory in any essential way, except that now the integration of the RG equation $\int_{1/L}^{1/d} K^{-1} dq/q$ gives $\ln^{1/2} L/d$ instead of conventional logarithmic dependence. Thus we find for the charged superconductor another law

$$\ln \Delta_{AB}/E_{AB} \sim -\frac{d}{\kappa_p a^2} \ln^{1/2} L/d . \quad (64)$$

Of course, to observe these dependences experimentally the ratio d/a must be not too large. Bearing in mind that we assumed $d > \xi$, we suggest superconductors with short coherence length (e.g., HTSC), or else $^4\text{He-II}$, although we are not aware of any experimental technique for preparing such narrow ($d \sim 20 \text{ \AA}$) long tubes for the helium experiment.

In this paper we mostly discussed the properties of SC in a sample with the ring topology threaded by the flux. We realize, however, that this situation can be modeled in a 1D wire

connected to the constant-current source through the Josephson contacts. In a way, here one deals with a reverse problem of finding the phase difference between the contacts as a function of current. This experimental setup might be of particular interest for studying the inelastic relaxation of SC due to their coupling to the substrate at large values of the current $j > j_c$, where j_c is the maximum value of the persistent current in Eq.(58). At mesoscopic values of the current $j \sim 1/L$, the phase slippage process in a 1D system is possible only if assisted by the energy dissipation to the substrate because we are dealing with the lowest energy levels, and 1D phonons form a discrete spectrum which is not in resonance with SC in the general case.

VI. ACKNOWLEDGMENTS

We would like to thank Yu.Kagan and I. Affleck for the discussions and valuable comments.

This work was supported by International Science Foundation (Grant No. MAA300) (NVP and BVS) and by Russian Foundation for Basic Research (Grants No. 94-02-05755a (VAK and AIP) and No. 95-02-06191a (NVP and BVS)), and partially by Grants No. INTAS-93-2834 [of the European Community] (BVS) and No. NWO-07-30-002 [of the Dutch Organization for Scientific Research] (NVP and BVS).

FIGURES

Fig.1. Spectrum evolution of a disordered system with increasing disorder. $N_a = 11$, number of particles $N_b = 7$, $U = 2.5$. Here $(2, 0)$ and $(0, 2)$ are the states with two identical minimal-momentum phonons; $(1, 1)$ is the state with two minimal-momentum phonons, moving in the opposite directions.

Fig.2. Spectrum evolution of a disordered system with increasing interaction. $N_a = 11$, $N_b = 7$, $W = 0.5$. The notation is the same as in Fig. 1.

Fig.3. Spectrum of the commensurate hard-core model as a function of the nearest-neighbor interaction; $N_a = 22$, $N_b = N_a/2$.

Fig.4. Scaling of relative AB-splitting with N_a in the case of single impurity ($\epsilon_1 = 0.5$); $N_a = 8, 10, 12, 14, 16, 18$; $N_b = N_a/2$.

Fig.5. Scaling of relative AB-splitting with N_a in the disordered case ($W = 0.5$); $N_a = 8, 10, 12, 14, 16, 18$; $N_b = N_a/2$. Each point represents an average over 100 realizations.

Fig.6. Scaling of the relative splitting of the first SC with N_a in the commensurate case; $N_a = 10, 12, 14, 16, 18, 20, 22$; $N_b = N_a/2$.

Fig.7. Scaling of the relative splitting of the first SC with N_a in the commensurate case at the critical point. $N_a = 10, 12, 14, 16, 18, 20, 22$, $N_b = N_a/2$.

Fig.8. The ratio R between the first SC energy splitting Δ_1 and the square of the lowest splitting Δ_{AB} ; $N_a = 11$, $N_b = 7$, and $W = 0.25$.

REFERENCES

- [1] B.V.Svistunov, J.Moscow Phys. Soc. **2**, 283 (1992).
- [2] Haldane F.D.M., Phys. Rev. Lett., **47**, 1840 (1981).
- [3] Popov V.N., *Functional Integrals in Quantum Field Theory and Statistical Physics*, Dordrecht: Reidel, 1983.
- [4] Coleman S., Phys. Rev., **D11**, 2088 (1975); Chui S.T. and P.A. Lee, Phys. Rev. Lett., **35**, 315 (1975).
- [5] Giamarchi T. and H.J.Schulz, Phys. Rev. B **37**, 325 (1988).
- [6] Kosterlitz J.M. and D.J.Thouless, J. Phys. C **6**, 1181 (1973).
- [7] Kosterlitz J.M., J. Phys. C **7**, 1046 (1974).
- [8] Kane C.L. and Fisher M.P.A., Phys. Rev. Lett. **68**, 1220 (1992). K.A.Matveev, D.Yue and L.I.Glazman, "Conduction of a Weakly Interacting 1D Electron Gas through a Single Barrier", preprint (1993).
- [9] Haldane F.D.M., J. Phys. C, **14**, 2585 (1981).
- [10] Luther A. and Peschel I., Phys. Rev. **B9** , 2911 (1974).
- [11] Mattis D.C., J. Math. Phys., **15**, 609 (1974).
- [12] Caldeira A. O. and Leggett A. J., Ann. Phys. **149**, 374 (1983).
- [13] Gogolin A.O. and Prokof'ev N.V., Phys. Rev., **B50**, 4921 (1994).
- [14] Leggett A. J. , Chakravarty S., Dorsey A. T., Fisher M. P. A., Garg A., and Zwerger W., Rev. Mod. Phys. **59**, 1 (1987) and refs. therein.
- [15] A.Schmid, Phys. Rev. Lett.**51**, 1506 (1983).
- [16] Kagan Yu., and Maksimov L.A., JETP **52**, 688 (1980).

- [17] Glazman L.I., Ruzin I.M., and Shklovskii B.I., Phys. Rev., **B45**, 8454 (1992).
- [18] Prokof'ev N.V., Int. J. Mod. Phys. **B 7**, 3327 (1993).
- [19] E.Dagotto, Rev. Mod. Phys., **66**, 763 (1994).
- [20] S.Pissanetzky, Academic, Orlando, 1984, ch.6.
- [21] B.Sutherland, Phys.Rev.Lett. **74**, 816 (1995).
- [22] V.A.Kashurnikov, A.I.Podlivaev, and B.V.Svistunov, Pis'ma v Z.Eksp.Teor.Fiz. **61**, 375 (1995).
- [23] I.Affleck, D.Gepner, H.J.Schulz, and T.Ziman, J.Phys. A: Math. Gen. **22**, 511 (1989).
And Refs. therein.
- [24] V.A.Kashurnikov and B.V.Svistunov (to be published).
- [25] The classification of levels adopted in [23] is based on the extra symmetry characteristic of the isotropic Heisenberg Hamiltonian and needs, generally speaking, some translation into the superfluid notation to make it possible to compare the results with the general theory of 1D superfluid-insulator transition. For one thing, the upper state of the first SC doublet is referred to in [23] as the lowest singlet excitation and the lower state of the SC doublet corresponds to the lowest triplet excitation of [23] (the one with zero z -projection of the total spin).
- [26] Meirav U., M.A. Kastner, M. Heiblum, and S.J. Wind, Phys. Rev. **B40**, 5871 (1989).
- [27] Chandrasekhar V., R.A.Webb, M.J.Brady, M.B.Ketchen, W.J.Galager and A.Kleinsasser, Phys. Rev. Lett. **67**, 3578 (1991); Maily D., Chapolier C., and Benoit A., *ibid* **70**, 2020 (1993).
- [28] Fazio R., F.W.J. Hekking, and A.A. Odintsov, Phys. Rev. Lett., **74**, 1843 (1995).

- [29] Hunklinger S., and Raychaudhuri A.K., in *Progress in Low Temperature Physics* , Vol.**X**, ed. Brewer D.F. (North-Holland, Amsterdam), ch.3 (1986).
- [30] Wen X.G., Phys. Rev, **B43**, 11025 (1990); Phys. Rev. Lett., **64**, 2206 (1990); Phys. Rev, **B44**, 5708 (1991).
- [31] Beenakker C.W.J., Phys. Rev. Lett., **64**, 216 (1990).
- [32] MacDonald A.H., Phys. Rev. Lett., **64**, 220 (1990).
- [33] Stone M., Phys. Rev., **42**, 8399 (1990).
- [34] Anderson P.W., Phys. Rev. **110**, 827 (1958); *ibid* **112**, 1900 (1958).